

Nonlinear Breathing-like Localized Modes in C₆₀ Nanocrystals

Alexander V. Savin^{1,2} and Yuri S. Kivshar²

¹*Semenov Institute of Chemical Physics, Russian Academy of Sciences, Moscow 119991, Russia*

²*Nonlinear Physics Center, Research School of Physics and Engineering,
Australian National University, Canberra ACT 0200, Australia*

We study the dynamics of nanocrystals composed of C₆₀ fullerene molecules. We demonstrate that such structures can support long-lived strongly localized nonlinear oscillatory modes, which resemble discrete breathers in simple lattices. We reveal that at room temperatures the lifetime of such nonlinear localized modes may exceed tens of picoseconds; this suggests that C₆₀ nanoclusters should demonstrate anomalously slow thermal relaxation when the temperature gradient decays in accord to a power law, thus violating the Cattaneo-Vernotte law of thermal conductivity.

I. INTRODUCTION

The fullerene era was started in 1985 with the discovery of a stable C₆₀ cluster and its interpretation as a cage structure with a soccer ball shape [1]. By now, many structural, electronic, and vibrational properties of fullerenes have been studied in detail. In spite of the rapidly increasing interest in new forms of fullerenes, icosahedral C₆₀, the "most beautiful molecule" [2], remains at the focus of active research as a prototype fullerene system. Fullerene molecule C₆₀ keeps a prominent position primarily for the high symmetry which greatly facilitates theoretical computations and the interpretation of experimental data, but also because it is only fullerene which can be grown in the form of large and nearly perfect single crystals. At room temperature, the diffraction experiments [3] show that the C₆₀ molecules are centered on sites of a face-centered-cubic (fcc) Bravais lattice. Each C₆₀ molecule in the solid-state phase is coupled to other C₆₀ molecules through weak Van-der Waals forces.

By now, the internal oscillations of the fullerene molecule C₆₀ are well studied, for the case of an isolated molecule and also for C₆₀ crystalline structures. Various theoretical and numerical methods have been developed and employed. However, the dynamics of fullerenes in the nonlinear regime and a role of nonlinear localized modes in such structures remain largely unexplored.

The main aim of this paper is to study strongly localized nonlinear oscillatory modes in such nanocrystals which resemble the discrete breathers of much simpler nonlinear lattices.

High symmetry of C₆₀ molecules allows the existence of internal nonlinear oscillations which do not alter substantially the spherical shape of the fullerene molecule. Such oscillations are excited by rotational degrees of freedom of pentagons and hexagons of the structure, with a variation of valent bonds. Resonant coupling of such high-amplitude oscillations with the neighboring C₆₀ molecules will remain weak, because due to conservation of a spherical shape the molecules will interact as large particle through Lennard-Jones potentials. This observation suggests that such structures can provide an ideal environment for the existence of nonlinear modes

strongly localized in the space [4, 5].

Here we demonstrate that the C₆₀ crystals can support long-lived strongly localized nonlinear modes which closely resemble the so-called discrete breathers in simple nonlinear lattices, and affect dramatically the heat relaxation of composite fullerene structures that obeys a power law instead of an exponential law.

II. MODEL AND SIMULATION METHODS

To analyze the dynamics of localized nonlinear modes in three-dimensional C₆₀ crystals, we consider a cubic nanocluster of C₆₀ molecules, as shown in Fig. 1 (a).

Inside of each C₆₀ molecule of the nanocluster two neighboring carbon atoms create a valent bond. We describe the energy associated with deformation of the valent bond created by two carbon atoms by the following interaction potential

$$U_1(\rho) = \epsilon_1 \{ \exp[-\alpha_0(\rho - \rho_0)] - 1 \}^2, \quad (1)$$

where ρ is the length of the valent bond, $\epsilon_1 = 4.9632$ eV and $\rho_0 = 1.418$ Å is the energy and the equilibrium length of the bond, parameter $\alpha_0 = 1.7889$ Å⁻¹. Each carbon atom is placed at the vertex of three planar valent angles. The corresponding deformation energy of a plane valent angle created by three atoms can be described by the interaction potential of the form,

$$U_2(\varphi) = \epsilon_2 (\cos \varphi + 1/2)^2, \quad (2)$$

where φ is value of the valent angle (equilibrium value of the angle $\varphi_0 = 2\pi/3$), energy $\epsilon_2 = 1.3143$ eV. Each valent bond is simultaneously belong to two planes, so that the deformation energy of the angle formed by two such planes can be described by the following interaction potential,

$$U_i(\phi) = \epsilon_i (1 - \cos \phi), \quad (3)$$

where ϕ is the corresponding angle (in equilibrium, $\phi = 0$), and the index $i = 3, 4, 5$ describe the type of the valent angle. Energy $\epsilon_3 = \epsilon_4 = 0.499$ eV, $\epsilon_5 \ll \epsilon_4$, so




FIG. 1: (a) General view of a cubic nanocluster composed of $5 \times 5 \times 5$ fullerene molecules C_{60} (buckyballs $\text{C}_{60}\text{-Ih}$), (b) isolated molecule C_{60} and (c) dimensionless density of its frequency spectrum for thermal vibrations at $T = 3\text{K}$.

that the latter contribution to the total energy can be neglected.

More detailed discussion and motivation of our choice of the interaction potentials U_1, \dots, U_4 can be found in Ref. [6]. Such potentials have been employed for modeling of thermal conductivity of carbon nanotubes [7, 8] graphene nanoribbons [6, 10] and also in the analysis of their oscillatory modes [9, 11, 12].

We describe the interaction of carbon atoms belonging to different C_{60} molecules by the Lennard-Jones 12-6 potential

$$U_{LJ}(r) = 4\epsilon_{LJ}[(\sigma/r)^{12} - (\sigma/r)^6], \quad (4)$$

where r – distance between the atom centers, energy $\epsilon_{LJ} = 0.004556$ eV, and parameter $\sigma = 3.851$ Å.

To verify our results, we also employ the simpler model where all interacting potentials between the atoms are described, instead of the set of the potentials U_1, \dots , and U_4 , by the Brenner potential [13].

III. INTRAMOLECULAR MODES

The high symmetry of an isolated, isotopically pure C_{60} molecule in its equilibrium configuration imposes very strong constraints on the form of the normal mode displacement patterns. All 60 sites are symmetry-equivalent, and the point group is the full icosahedral group I_h . Although there are 180

degrees of freedom (3×60) for each C_{60} molecule, the icosahedral symmetry gives rise to a number of degenerate modes, so that only 46 distinct mode frequencies are expected for the C_{60} molecule. All vibrational modes of C_{60} are distributed [14, 15] as $2A_g + A_u + 4T_{1g} + 5T_{1u} + 4T_{3g} + 5T_{3u} + 6G_g + 6G_u + 8H_g + 7H_u$.

Dynamics of an isolated C_{60} fullerene molecule is described by a system of 3×60 nonlinear equations of the standard form, $M\ddot{\mathbf{u}} = -\partial H / \partial \mathbf{u}$, where $M = 12 \cdot 1.6603 \cdot 10^{-27}$ kg – mass of the carbon atom, system Hamiltonian is $H = M(\dot{\mathbf{u}}, \dot{\mathbf{u}})/2 + E(\mathbf{u})$, and 180-dimensional vector $\mathbf{u} = \{(x_n, y_n, z_n)\}_{n=1}^{60}$ defines the coordinates of all atoms of the complex molecule. Potential energy of the molecule $E(\mathbf{u})$ is composed by all types of interactions described by the potentials U_1, \dots, U_4 discussed above.

To find the equilibrium stationary state of the C_{60} molecule, we need to solve numerically the minimization problem for the energy functional, $E(\mathbf{u}) \rightarrow \min$. In the model under consideration, the molecule C_{60} is described by the set of coordinates \mathbf{u}_0 , and it has a shape of perfect icosahedron [see Fig. 1 (b)] with the radius $R = 3.514$ Å, for which all valent bonds have the equilibrium length $\rho = \rho_0$.

Linearizing the model near the equilibrium stationary state, we obtain a system of linear equations $M\ddot{\mathbf{u}} = \mathbf{A}\mathbf{u}$, where \mathbf{A} is 180×180 dimensional matrix, $\mathbf{A} = \{\partial^2 E / \partial u_i \partial u_j |_{\mathbf{u}=\mathbf{u}_0}\}_{i,j=1}^{180}$. Diagonalizing the matrix \mathbf{A} we find all 180 eigenmodes of the molecule. The first six zero-frequency modes correspond to rigid translations

TABLE I: Values ω (in cm^{-1}) for the 46 distinct normal mode frequencies of an isolated C_{60} molecule (n is mode degeneracy). Notations of the modes are from Ref. [15].

mode	ω	n	mode	ω	n	mode	ω	n	mode	ω	n
$H_g(1)$	162.1	5	$H_g(3)$	497.4	5	A_u	776.0	1	$G_g(5)$	1437.5	4
$T_{3u}(1)$	223.2	3	$T_{3u}(2)$	497.8	3	$G_u(4)$	847.7	4	$T_{1g}(3)$	1454.9	3
$G_u(1)$	225.5	4	$H_u(4)$	501.8	5	$T_{3u}(3)$	890.2	3	$A_g(2)$	1503.2	1
$H_u(1)$	278.2	5	$T_{1u}(2)$	503.6	3	$H_g(5)$	1020.0	5	$T_{3u}(5)$	1511.7	3
$H_g(2)$	282.8	5	$G_g(3)$	534.7	4	$G_g(4)$	1039.8	4	$T_{1u}(4)$	1512.9	3
$G_g(1)$	323.1	4	$G_u(2)$	535.0	4	$H_u(5)$	1121.8	5	$H_g(7)$	1518.2	5
$H_u(2)$	358.6	5	$T_{3g}(2)$	542.5	3	$T_{1u}(3)$	1181.2	3	$G_u(6)$	1540.6	4
$T_{1u}(1)$	364.3	3	$H_u(3)$	557.1	5	$T_{3g}(4)$	1267.6	3	$G_g(6)$	1553.8	4
$T_{1g}(1)$	398.5	3	$G_u(3)$	579.3	4	$H_g(6)$	1270.3	5	$H_u(7)$	1575.0	5
$G_g(2)$	413.1	4	$T_{3g}(3)$	603.1	3	$T_{3u}(4)$	1272.0	3	$H_g(8)$	1579.3	5
$A_g(1)$	418.6	1	$H_g(4)$	673.7	5	$H_u(6)$	1376.2	5			
$T_{3g}(1)$	442.3	3	$T_{1g}(2)$	675.7	3	$G_u(5)$	1399.8	4			

and rotations of the molecule. Other 174 modes are generated and, in view of their degeneracy, there are exists 46 distinct modes – see Table I.

The frequencies of all oscillatory eigenmodes of an isolated C_{60} molecule are summarized in Table I. All frequency are located in the interval $162.1 \leq \omega \leq 1579.3 \text{ cm}^{-1}$. Definitely, the model with only 6 parameters is not able to predict exactly the frequencies obtained in the framework of the first principles calculations or experimental results, see [15]. Nevertheless, this model demonstrates a correct structure of the eigenmode frequencies and predict rather well the frequency band measured experimentally, $272 \leq \omega \leq 1575 \text{ cm}^{-1}$.

It is important to mention that the use of the Brenner potentials [13] instead of the set of the potentials U_1, \dots, U_4 leads much broader frequency band, $182.9 \leq \omega \leq 1738.9 \text{ cm}^{-1}$. In addition, the Brenner potential do not predict correctly the degeneracy of the frequency spectrum, so that the equilibrium shape of the C_{60} molecule deviates from the icosahedra shape.

In order to model thermal oscillations of the molecule, we start from its equilibrium state and excite each of its eigenmode with the energy $k_B T$. The frequency spectrum of the C_{60} molecule at the temperature $T = 3\text{K}$ is shown in the Fig. 1 (c). As expected, the spectrum is discrete, and the number of peaks coincides with the number of eigenmodes, whereas the peak positions coincide with the corresponding eigenfrequencies. At low temperatures, the oscillations are linear, so that the spectrum corresponds to a system of uncoupled linear oscillators.

If we increase the temperature to $T = 30\text{K}$, the discrete nature of the spectrum remains unchanged, but high-frequency modes demonstrate some energy exchange and interaction. For much higher temperatures such as $T = 300\text{K}$ the discrete nature of the spectrum is lost and the oscillations become strongly nonlinear demonstrating interaction with strong energy exchange.

IV. LOCALIZED NONLINEAR MODES

The C_{60} molecules occupy fcc sites in the solid phase, and at room temperature, the rapid rotational diffusion of each molecule leads to an effective fcc crystal structure. To study localized modes of the fullerene molecule placed in a crystal, we consider a cubic nanocluster composed of $5 \times 5 \times 5$ molecules C_{60} , as shown in Fig. 1 (a).

First, we find the ground state of the nanocluster by minimizing its energy. Next, for the central molecule we excite its n -th eigenmode ($n = 7, \dots, 180$) with the energy E_0 and study the decay of the kinetic energy in the cluster. Due to the nonvalent coupling between the fullerenes, the energy is transferred from one molecule to another. To model the energy spreading in an effectively infinite crystal, we introduce lossy boundary conditions, when all surface molecules experience damping with the relaxation time $t_r = 20 \text{ ps}$. If the excitation of the certain eigenmode lead to the creation of a localized state, or breathing-like mode, the kinetic energy will not vanish but instead will approach a certain nonzero value. Otherwise, the initial excitation will vanish completely, so that the kinetic energy of the central molecule $E_k \rightarrow 0$ when $t \rightarrow \infty$.

First, we excite with the energy $E_0 = 0.2$ the n -th mode ($n = 7, \dots, 180$) of the central molecule of the fullerene crystal. Depending on the type of the modes, our numerical results reveal three different scenarios of the mode relaxation, as shown in Fig. 2 (a-c). The first scenario is the rapid energy relaxation and decay of the mode oscillations (see curves 1, 4, and 7); the second scenario is slow relaxation (curves 2, 5, and 8), and the third scenario is the generation of stationary spatially localized nonlinear modes (curves 3, 6, and 9).

The first and second scenarios are observed for the modes associated with a periodic change of the spherical shape or the size of the C_{60} molecule. The most interesting third scenario is observed for the modes associated




FIG. 2: Evolution of the kinetic energy E_k of the central fullerene C_{60} molecule in nanocluster composed of $5 \times 5 \times 5$ molecules packed in the form of a fcc cubic crystal. At the initial moment $t = 0$ only one eigenmode is excited for the central molecule with the energy $E_0 = 0.2$ eV. (a) curve 1 shows the decay of the mode $H_g(1)$ (frequency $\omega = 162.2 \text{ cm}^{-1}$), curve 2 – the mode $H_u(4)$ ($\omega = 501.8 \text{ cm}^{-1}$), and the curve 3 – the mode $T_{3g}(1)$ ($\omega = 442.3 \text{ cm}^{-1}$); (b) curve 4 gives the dependence for the mode $A_g(1)$ ($\omega = 418.6 \text{ cm}^{-1}$), curve 5 – the mode $T_{3g}(4)$ ($\omega = 1267.6 \text{ cm}^{-1}$), curve 6 – the mode $T_{1g}(2)$ ($\omega = 675.7 \text{ cm}^{-1}$); (c) curve 7 describes the decay of the mode A_u (frequency $\omega = 776.0 \text{ cm}^{-1}$), curve 8 – the mode $G_g(5)$ ($\omega = 1437.5 \text{ cm}^{-1}$), and the curve 9 – the mode $T_{1g}(3)$ ($\omega = 1454.9 \text{ cm}^{-1}$).

with the motion of atoms along the spherical surface so that the molecule keeps its form almost unchanged. These are the eigenmodes $T_{3g}(1)$, $T_{1g}(2)$, and $T_{1g}(3)$ presented in Fig. 3. For all those modes, the molecule conserves its spherical shape and therefore the resonant interatomic interaction is reduced.

Figure 4 shows the spatial distribution of the kinetic energy of the nonlinear localized state generated by the excitation of the eigenmode $T_{3g}(1)$. The main part of the energy is concentrated at the central C_{60} molecule, so this state resembles discrete breathers studied earlier for simpler lattices. The nonlinear states generated by the eigenmodes $T_{1g}(2)$ $T_{1g}(3)$ are localized even stronger. For $E > 0.01$ eV, all such modes represent strongly anharmonic oscillations with the frequency decaying with




FIG. 3: Selected eigenmodes of an isolated C_{60} molecule which lead to the generation of spatially localized nonlinear modes (breathers) of the fullerene crystal. (a) mode $T_{3g}(1)$ (frequency $\omega = 442.3 \text{ cm}^{-1}$); (b) mode $T_{1g}(2)$ (frequency $\omega = 675.7 \text{ cm}^{-1}$); (c) mode $T_{1g}(3)$ (frequency $\omega = 1454.9 \text{ cm}^{-1}$).




FIG. 4: Spatial profile of the kinetic energy E of the breathing-like nonlinear localized mode of the central fullerene molecule in the nanocluster of $5 \times 5 \times 5$ of C_{60} molecules in the planes (a) xy and (b) xz , intersecting the central molecule (shown by a black circle). For the generation of such breathers, the eigenmode $T_{3g}(1)$ was excited with the energy $E = 0.2$ eV.

the mode amplitude. The mode $T_{1g}(2)$ carries the largest amount of energy that can reach 1 eV.

It is important to mention that the model employing the Brenner potentials predicts the existence of a larger number of localized modes.

In a thermalized nanocrystal all localized modes have a finite life time. Their decay occurs mainly due to the interaction with other modes of the same molecule. Therefore, in order to define the lifetime of a nonlinear mode, it would be sufficient to study the dynamics of an isolated C_{60} molecule with one strongly excited mode corresponding to the nonlinear localized state and all other modes weakly thermalized.

In Fig. 5 we show the interaction of one large-amplitude eigenmode $T_{3g}(1)$ with all other weakly thermalized eigenmodes of the fullerene C_{60} molecule. As follows from those results, the energy of the localized mode decays monotonically exciting other modes. When the initial energy of the $T_{3g}(1)$ mode is $E_1 = 0.2$ eV and the energy of all other modes is $E_0 = 0.002$ eV, the localized mode loses a half of its energy for $t_0 = 1.4$ ns, for $E_0 = 0.004$ eV – this time is $t_0 = 0.8$ ns, for $E_0 = 0.008$ eV – $t_0 = 0.4$ ns, and for $E_0 = 0.016$ eV – $t_0 = 0.1$ ns. Highly-excited mode $T_{1g}(2)$ with the energy $E_1 = 0.2$ eV loses a half of its energy for $t_0 = 0.8$ ns if energy $E_0 = 0.002$ eV,




FIG. 5: Interaction of a single large-amplitude mode T3g(1) with all other weakly thermalized eigenmodes of C₆₀ molecules. As the initial condition at $t = 0$ we excite the eigenmode T3g(1) with the frequency $\omega = 442.3 \text{ cm}^{-1}$ and the energy $E_1 = 0.2 \text{ eV}$, and all other eigenmodes with the energy $E_0 = 0.002 \text{ eV}$. Shown is the evolution of the energy distribution over the frequencies. The peak corresponds to a large-amplitude mode (shown in red).

for $t_0 = 0.5 \text{ ns}$ if $E_0 = 0.004 \text{ eV}$, for $t_0 = 0.3 \text{ ns}$ if $E_0 = 0.008 \text{ eV}$, and for $t_0 = 0.1 \text{ ns}$ if $E_0 = 0.016 \text{ eV}$. The smallest lifetime is shown by the eigenmode T_{1g}(3), which for the input energy $E_1 = 0.2 \text{ eV}$ and $E_0 = 0.002$ loses a half of its energy for $t_0 = 0.1 \text{ ns}$.

Therefore, our numerical results demonstrate that in a crystal formed by C₆₀ molecules the spatially localized nonlinear modes are long-lived excitations, and at room temperature their life time is of the order of several picoseconds.

V. THERMAL RELAXATION OF C₆₀ CRYSTAL

The existence of long-lived nonlinear localized collective modes in the nanocluster of C₆₀ fullerene molecules should affect substantially their physical properties. If the thermal energy can be concentrated for long time in some localized spots, such a crystal is expected to demonstrate low thermal conductivity. To confirm this idea, we consider thermal relaxation of a cubic nanocluster composed of $10 \times 12 \times 12$ C₆₀ molecules, as shown in Fig. 6.

The 180-dimensional vector \mathbf{u}_{nmk} define the coordinates of the atoms of (n, m, k) -th molecule of the nanocluster ($n = 1, \dots, 10$; $m = 1, \dots, 12$; $k = 1, \dots, 12$). For thermalizing the nanocluster, we solve the system of Langevin equations

$$M\ddot{\mathbf{u}}_{nmk} = -\partial H / \partial \mathbf{u}_{nmk} - \Gamma M \dot{\mathbf{u}}_{nmk} + \Xi_{nmk}, \quad (5)$$

where $\Gamma = 1/t_r$ is the Langevin collision frequency with $t_r = 0.1 \text{ ps}$ being the corresponding particle relaxation time, and $\Xi_{nmk} = \{\xi_{nmk,i}\}_{i=1}^{180}$ is a 180-dimensional vector of Gaussian distributed stochastic forces describing the interaction of atom with the thermostat with correlation functions $\langle \xi_{n_1 m_1 k_1, i}(t_1) \xi_{n_2 m_2 k_2, j}(t_2) \rangle =$




FIG. 6: Relaxation the thermal gradient ΔT between the opposite sides of the cubic cluster of $10 \times 12 \times 12$ fullerene C₆₀ molecules with time t . Left insert shows this dependence in the log-scale, with a straight line corresponding to a power-law dependence, $\Delta T \sim t^{-1}$. Right insert shows a cubic nanocluster with an arrow indicating the heat flow. Dependencies 1 and 4 correspond to the approximation of rigid molecules when their internal dynamics is neglected, whereas curves 2 and 5 show the results of the full molecular dynamics simulations. Curves 3 and 6 show the results for the model employing the Brenner potentials.

$2M\Gamma k_B T_{n_1 m_1 k_1} \delta_{n_1 n_2} \delta_{m_1 m_2} \delta_{k_1 k_2} \delta_{ij} \delta(t_2 - t_1)$, T_{nmk} is the temperature of the molecule with the site (n, m, k) .

To obtain the initial thermalized state of the nanocluster we solve numerically the Langevin equations (5) with the initial conditions corresponding to the equilibrium state. We run the simulations for $t = 10t_r$, and then switch off the coupling with the thermostat, i.e. neglecting the last two terms in Eqs. (5), in order to study the thermal relaxation of the nanocluster.

We thermalize the nanocluster asymmetrically, to have only one side heated to the temperature $T = 500 \text{ K}$, whereas keeping all other atoms at the temperature $T = 300 \text{ K}$. To achieve this, we solve the system of Langevin equations at $T_{nmk} = 500 \text{ K}$ for the atoms with the index $k = 1$ but for all other atoms with the indices $k = 2, \dots, 12$ we put $T_{nmk} = 300 \text{ K}$. In this case, we achieve a temperature different of $\Delta T = 200 \text{ K}$ between the opposite sides of the C₆₀ nanocube. As the next step, we study the thermal relaxation of the temperature gradient.

As a result of thermal energy exchange between the fullerene molecules, temperature difference should decay monotonically. The crystal of fullerene molecules C₆₀ has a three-dimensional structure so it should obey the Fourier law for thermal conductivity. Because time does not enter the Fourier law, it does not allow to determine the temporal relaxation of the thermal gradient, for which we should employ the well-know generalization known as Cattaneo-Vernotte (CV) law [16, 17]. In its one-dimensional version, this law can be formulated through the differential equations,

$$\left(1 + \tau \frac{\partial}{\partial t}\right) \vec{q} = -\kappa \vec{\nabla} T, \quad (6)$$

where κ is the standard coefficient of the thermal conductivity and τ is the characteristic relaxation time of the system. In according to Eq. (6), the temperature gradient ΔT should decay in accord with an exponential dependence.

Dependence of the temperature gradient ΔT vs. t is shown in Fig. 6. As follows from this dependence, the temperature difference decays very slow, in accord with the power-law dependence, so that it does not obey the Cattaneo-Vernotte law. In order to understand the reason for such an anomalous thermal relaxation as well as the role played by the internal dynamics of the molecules, we replace the C_{60} molecules by rigid balls and solve the same problem assuming that the molecules in the crystal interact between each other as structureless particles. Our numerical results show that in this approximation the relaxation of the temperature difference ΔT follows an exponential dependence, in accord with the Cattaneo-Vernotte law. Therefore, the anomalously slow relaxation can be attributed to the excitation of long-lived nonlinear oscillations of the C_{60} nanocluster.

VI. CONCLUSIONS

We have applied the molecular-dynamics numerical simulations to study the nonlinear dynamics and ther-

mal relaxation of crystalline structures composed of C_{60} fullerene molecules. We have revealed that such complex nanoclusters support a special type of strongly localized nonlinear modes resembling discrete breathers of simple nonlinear lattices. For such modes, the kinetic energy is localized primarily in the rotational modes of a single C_{60} molecule and it decays slowly inside the fullerene nanocrystal. Importantly, our results have demonstrated that at room temperatures the lifetime of such nonlinear localized modes may exceed tens of picoseconds. The existence of such long-lived localized nonlinear states in the nanoclusters explains the anomalously slow thermal relaxation observed in such structures when the temperature gradient decays in accord with the power but not exponential law, thus violating the Cattaneo-Vernotte law of thermal conductivity.

Acknowledgements

Alex Savin acknowledges a warm hospitality of the Nonlinear Physics Center at the Australian National University, and he thanks the Joint Supercomputer Center of the Russian Academy of Sciences for the use of their computer facilities. The work was supported by the Australian Research Council.

-
- [1] H. W. Kroto, J.R. Heath, S. C. O'Brien, R. F. Curl, R. E. Smalley, *Nature* **318**, 162 (1985).
 - [2] H. Aldersey-Williams: *The Most Beautiful Molecule: The Discovery of the Buckyball*, (Wiley, New York, 1995).
 - [3] R. M. Fleming, T. Siegrist, P. M. Marsh, B. Hessen, A. R. Kortan, D. W. Murphy, R. C. Haddon, R. Tycko, G. Dabbagh, A. M. Muzice, M. L. Kaplan, and S. M. Zahurak, *Mater. Res. Soc. Symp. Proc.* **206**, 691 (1991).
 - [4] D. K. Campbell, S. Flach, and Yu. S. Kivshar, *Phys. Today* **57**, 43 (2004).
 - [5] S. Flach and A. V. Gorbach, *Phys. Rep.* **467**, 1 (2008).
 - [6] A. V. Savin, Yu. S. Kivshar, and B. Hu, *Phys. Rev. B* **82**, 195422 (2010).
 - [7] A. V. Savin, B. Hu, and Yu. S. Kivshar, *Phys. Rev. B* **80**, 195423 (2009).
 - [8] A. V. Savin, Yu. S. Kivshar, and B. Hu, *EPL*, **88**, 26004 (2009).
 - [9] A. V. Savin and Yu. S. Kivshar, *Appl. Phys. Lett.* **94**, 111903 (2009).
 - [10] A. V. Savin, Yu. S. Kivshar. *Letters on Materials* **1**, 3 (2011). (in Russian)
 - [11] A. V. Savin and Yu. S. Kivshar, *Europhys. Lett.* **89**, 46001 (2010).
 - [12] A. V. Savin and Yu. S. Kivshar, *Phys. Rev. B* **81**, 165418 (2010).
 - [13] D. W. Brenner, O. A. Shenderova, J. A. Harrison, S. J. Stuart, B. Ni, and S. B. Sinnott, *J. Phys.: Condens. Matter* **14**, 783 (2002).
 - [14] D. E. Weeks, W. G. Harter, *J. Chem. Phys.* **90**, 4744 (1989).
 - [15] J. Menendez and J. B. Page, In: *Light Scattering in Solids VIII*, Eds. M. Cardona and G. Guntherodt (Springer-Verlag, Berlin, 2000).
 - [16] P. Vernotte, C. R. Hebd, *Seances Acad. Sci.* **246**, 3154 (1958).
 - [17] C. Cattaneo, C. R. Hebd, *Seances Acad. Sci.* **247**, 431 (1958).